

Combining a Comprehensive Physical Therapy Program and Electroshock Therapy for a Patient with Plantar Fasciitis: A Case Report

Kathryn Piersiak, BS, ATC, Doctor of Physical Therapy Student
Doctor of Physical Therapy Program, University of New England, Portland, Maine

Background

- Plantar fasciitis (PF) affects 2 million Americans per year and 10% of the population over a lifetime.¹
- PF is inflammation to the thick, fibrous connective tissue originating on the medial calcaneal tubercle extending to the metatarsal heads of the foot.^{2,3}
- A comprehensive physical therapy (PT) plan of care (POC) of proximal strengthening, distal stretching and soft tissue massage (STM) has been shown to improve PF symptoms.
- Extracorporeal shockwave therapy (ESWT) is a non-invasive modality that uses sound waves to create a controlled microtrauma to the impaired tissue to stimulate a healing response and micro vascularization.⁴
- ESWT has shown to be effective in patients with PF, but has not been used in conjunction with a comprehensive PT program.

Purpose

The purpose of this case report was to investigate the recovery of a patient with PF while using a combined approach of conventional therapy, targeted hip strengthening, and electroshock therapy.

Case Description

- The patient was a 48-year-old female presenting with right plantar fasciitis.
- She underwent one treatment of ESWT two months prior to initial evaluation.
- She was scheduled for additional ESWT treatment after PT.
- The patient had a history of failed PT for the same injury last year.
- She reported pain as an aching/dull/throbbing sensation with sharp pains during ambulation
- Patient described difficulty with weight bearing activities including walking, stairs, running, and standing on her feet at work.




Figure 1: Plantar Fascia Anatomy⁵

Timeline

History of Treatment

- Prior physical therapy for 3 months
- ESWT done one month prior to initial evaluation
- Referred to physical therapy

Initial Evaluation

- Tenderness along medial and origin of plantar fascia, calcaneus, and along dorsal foot metatarsals 2-4
- Trendelenburg left hip

Visits 1-2

- Initiation of manual therapy
- Arch tape trial
- Initiation of stretching to patient tolerance
- Home exercise program (HEP) given to patient

Visits 3-4

- Progression of HEP
- Progression of therex to include more strengthening
- Talocrural joint mobilizations to improve range of motion (ROM)

Visits 5-6

- Introduction of intensive hip strengthening program
- Progression of digital distraction to improve ROM for gait mechanics

Visits 7-8

- Re-evaluation of outcome measures
- Repeat ECSW Therapy to be completed 1 month after final re-evaluation

Interventions




Figure 2: Therapeutic Interventions

Manual Therapy:

- Soft tissue massage to gastrocnemius/soleus
- Great toe extension stretch
- Talocrural anterior-posterior mobilizations grade 2-3
- Digital distraction

Therapeutic Exercise:

- Towel scrunches
- Ankle 4 way theraband
- Great toe stretch with ankle 90 degrees
- Runner stretch with bent knee
- Myofascial release stick to gastrocnemius, anterior tibialis, and hamstrings
- Marble pick-ups
- Lateral walking with theraband
- Single leg stance balance on Airex®
- Bilateral side lying clamshells

Modalities:

- GameReady®
- Ice foot roll

Arch Taping:

- Medial longitudinal arch tape





Figure 3: Medial Longitudinal Arch Taping

Outcomes



- Palpation tenderness was only present over PF insertion.
- The patient's gait no longer demonstrated a positive Trendelenburg on the left side upon re-evaluation.

Conclusion

- A comprehensive physical therapy program including targeted hip strengthening and ESWT therapy for a patient with PF resulted in increased functional abilities, increased strength, and a decrease in pain.
- While there is limited research about ESWT therapy for the treatment of PF, when combined with conventional therapy, there was a positive outcome.
- Additional research is necessary to determine the effectiveness of targeted hip strengthening and extracorporeal shockwave therapy specifically for chronic cases of PF.

Acknowledgements

- The author acknowledges Kirsten Buchanan, PhD, PT, ATC, for case report for both conceptualization and editing, and Marissa Valle, PT, DPT for assistance of patient supervision. The author also acknowledges the patient for their willingness to participate in this case report.
- Please address all correspondence to Kate Piersiak at: kathryn.piersiak@gmail.com.

References

- Martin RL, Davenport TE, Reischl SF, et al. Heel pain-plantar fasciitis: revision 2014. *J Orthop Sports Phys Ther.* 2014;44(11):A1-A23. doi: 10.2519/jospt.2014.0303
- Schwartz EN, Su J. Plantar fasciitis: a concise review. *Perm J.* 2014;18(1):e105-e107. doi:10.7812/TPP/13-113
- Sonu P, Aman. Physiotherapy treatment in plantar fasciitis: a case report. *Indian J Physiother and Occup Ther.* 2015;9(1):54-58. doi: 10.5958/0973-5674.2015.00012.X
- Kamoneki DH, Gonçalves GA, Yi LC, Júnior IL. Effect of stretching with and without muscle strengthening exercises for the foot and hip in patients with plantar fasciitis: a randomized controlled single-blind clinical trial. *Man Ther.* 2016;23:76-82. doi: //doi.org/10.1016/j.math.2015.10.006
- Plantar Fasciitis. Local Physio Website <https://www.local-physio.co.uk/articles/heel-pain/plantar-fasciitis/>. Updated 2019. Accessed November 22, 2019.